

Spaces not distinguishing various kinds of convergence of sequences of real functions

Rafał Filipów



37th International Summer Conference on Real Functions Theory
2023 Rowy, Poland

New results in my talk are from a joint work with Adam Kwela:

“Spaces not distinguishing ideal pointwise and σ -uniform convergence”

which is available at arXiv (2308.09557).

Redefinition

- Topological space = nonempty normal (a.k.a. T_4) space
- Function = continuous real-valued function defined on X ($f : X \rightarrow \mathbb{R}$)

Four kinds of convergence of sequences of functions

Pointwise convergence

$$(f_n) \xrightarrow{P} 0 \text{ if } \forall \varepsilon > 0 \forall x \in X \exists k \forall n > k (|f_n(x)| < \varepsilon)$$

Equal convergence (1979 – Császár-Laczkovich)

Quasi-normal convergence (1991 – Bukovská)

$$(f_n) \xrightarrow{QN} 0 \text{ if } \exists (\varepsilon_n) \rightarrow 0 \forall x \in X \exists k \forall n > k (|f_n(x)| < \varepsilon_n)$$

Uniform convergence

$$(f_n) \xrightarrow{U} 0 \text{ if } \forall \varepsilon > 0 \exists k \forall n > k \forall x \in X (|f_n(x)| < \varepsilon)$$

σ -uniform convergence

$$(f_n) \xrightarrow{\sigma-U} 0 \text{ if there is a partition } X_1, X_2, \dots \text{ of } X \text{ such that}$$
$$\forall k \in \mathbb{N} \quad (f_n \upharpoonright X_k) \xrightarrow{U} 0$$

Four kinds of convergence of sequences of functions

Relationships

Theorem (Easy)

$$(f_n) \xrightarrow{U} 0 \implies (f_n) \xrightarrow{\sigma-U} 0 \quad \text{and} \quad (f_n) \xrightarrow{QN} 0 \implies (f_n) \xrightarrow{P} 0$$

Theorem (Császár-Laczkovich, 1979)

$$(f_n) \xrightarrow{\sigma-U} 0 \iff (f_n) \xrightarrow{QN} 0$$

Question

$$(f_n) \xrightarrow{U} 0 \stackrel{?}{\iff} (f_n) \xrightarrow{\sigma-U} 0 \quad \text{and} \quad (f_n) \xrightarrow{QN} 0 \stackrel{?}{\iff} (f_n) \xrightarrow{P} 0$$

Theorem (Easy)


$$\forall (f_n: X \rightarrow \mathbb{R}) \left((f_n) \xrightarrow{U} 0 \iff (f_n) \xrightarrow{\sigma-U} 0 \right) \iff X \text{ is finite}$$

Definition (Bukovský-Reclaw-Repický, 1991)

$$X \text{ is a QN-space} \iff \forall (f_n: X \rightarrow \mathbb{R}) \left((f_n) \xrightarrow{QN} 0 \iff (f_n) \xrightarrow{P} 0 \right)$$

Four kinds of convergence of sequences of functions

Diagram of relationships



$$(f_n) \xrightarrow{U} 0 \implies (f_n) \xrightarrow{\sigma-U} 0, \dots$$

$$\forall (f_n: X \rightarrow \mathbb{R}) \left((f_n) \xrightarrow{U} 0 \iff (f_n) \xrightarrow{\sigma-U} 0 \right) \iff X \text{ is finite}$$

$$X \text{ is a QN-spaces} \iff \forall (f_n: X \rightarrow \mathbb{R}) \left((f_n) \xrightarrow{QN} 0 \iff (f_n) \xrightarrow{P} 0 \right)$$

Definition

A family $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$ is an **ideal** on \mathbb{N} if

- 1 $A \subseteq B \in \mathcal{I} \implies A \in \mathcal{I}$,
- 2 $A, B \in \mathcal{I} \implies A \cup B \in \mathcal{I}$,
- 3 \mathcal{I} contains all finite subsets of \mathbb{N} and $\mathbb{N} \notin \mathcal{I}$.

Example

- 1 $\text{Fin} = \{A \subseteq \mathbb{N} : A \text{ is finite}\}$
- 2 $\mathcal{I}_{1/n} = \left\{ A \subseteq \mathbb{N} : \sum_{n \in A} \frac{1}{n} < \infty \right\}$
- 3 $\mathcal{I}_d = \left\{ A \subseteq \mathbb{N} : \lim_{n \rightarrow \infty} \frac{|A \cap \{1, \dots, n\}|}{n} = 0 \right\}$

Ideal convergence of sequences of reals

Definition

A sequence (x_n) of reals is **convergent to zero** if

$$\forall \varepsilon > 0 \quad \exists k \forall n > k \quad (|x_n| < \varepsilon)$$

equivalently:

$$\forall \varepsilon > 0 \quad \exists A \in \text{Fin} \quad \forall n \in \mathbb{N} \setminus A \quad (|x_n| < \varepsilon)$$

Definition

Let \mathcal{I} be an ideal on \mathbb{N} .

A sequence (x_n) of reals is **\mathcal{I} -convergent to zero** if

$$\forall \varepsilon > 0 \quad \exists A \in \mathcal{I} \quad \forall n \in \mathbb{N} \setminus A \quad (|x_n| < \varepsilon).$$

—
In symbols: $(x_n) \xrightarrow{\mathcal{I}} 0$

Ideal convergence of sequences of reals

Examples

Definition

A sequence (x_n) of reals is **\mathcal{I} -convergent to zero** ($(x_n) \xrightarrow{\mathcal{I}} 0$) if

$$\forall \varepsilon > 0 \quad \exists A \in \mathcal{I} \quad \forall n \in \mathbb{N} \setminus A \quad (|x_n| < \varepsilon).$$

Example

Since $\text{Fin} \subseteq \mathcal{I}$,

$$\text{if } (x_n) \rightarrow 0, \text{ then } (x_n) \xrightarrow{\mathcal{I}} 0.$$

Example

Let $x_n = \begin{cases} 1 & \text{if } n = 1, 4, 9, \dots, k^2, \dots, \\ 0 & \text{otherwise.} \end{cases}$

Then (x_n) is a **divergent sequence**, but

$$(x_n) \xrightarrow{\mathcal{I}_{1/n}} 0,$$

because $\sum_{k=1}^{\infty} \frac{1}{k^2} < \infty$, so $A = \{1, 4, 9, \dots, k^2, \dots\} \in \mathcal{I}_{1/n}$.

Ideal convergence of sequences of functions

Let \mathcal{I} be an ideal on \mathbb{N} . Let $f_n : X \rightarrow \mathbb{R}$, $n = 1, 2, \dots$

Ideal pointwise convergence

$$(f_n) \xrightarrow{\mathcal{I}-P} 0 \text{ if } \forall_{\varepsilon > 0} \forall_{x \in X} \exists_{A \in \mathcal{I}} \forall_{n \in \mathbb{N} \setminus A} (|f_n(x)| < \varepsilon)$$

Ideal quasi-normal convergence

$$(f_n) \xrightarrow{\mathcal{I}-QN} f \text{ if } \exists_{(\varepsilon_n) \xrightarrow{\mathcal{I}} 0} \forall_{x \in X} \exists_{A \in \mathcal{I}} \forall_{n \in \mathbb{N} \setminus A} (|f_n(x)| < \varepsilon_n)$$

Ideal uniform convergence



$$(f_n) \xrightarrow{\mathcal{I}-U} 0 \text{ if } \forall_{\varepsilon > 0} \exists_{A \in \mathcal{I}} \forall_{n \in \mathbb{N} \setminus A} \forall_{x \in X} (|f_n(x)| < \varepsilon)$$

Ideal σ -uniform convergence


$$(f_n) \xrightarrow{\mathcal{I}-\sigma-U} 0 \text{ if there is a partition } X_1, X_2, \dots \text{ of } X \text{ such that}$$
$$\forall_{k \in \mathbb{N}} (f_n \upharpoonright X_k) \xrightarrow{\mathcal{I}-U} 0$$

Ideal convergence of sequences of functions

Diagram of relationships



Minimal size of a space which distinguishes convergence




Definition

- \bullet $\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-QN}) = \min\{|X| : \exists f_n: X \rightarrow \mathbb{R} (f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \wedge f_n \not\xrightarrow{\mathcal{I}\text{-QN}} 0)\}$
- \bullet $\text{non}(\mathcal{I}\text{-QN}, \mathcal{I}\text{-}\sigma\text{-U}) = \min\{|X| : \exists f_n: X \rightarrow \mathbb{R} (f_n \xrightarrow{\mathcal{I}\text{-QN}} 0 \wedge f_n \not\xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0)\}$
- \bullet $\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \min\{|X| : \exists f_n: X \rightarrow \mathbb{R} (f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \wedge f_n \not\xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0)\}$

Minimal size of a space which distinguishes convergence

Relationship



Theorem (Easy)

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \min\{\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-QN}), \text{non}(\mathcal{I}\text{-QN}, \mathcal{I}\text{-}\sigma\text{-U})\}$$

Proof

- If $|X| < \text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U})$, then \dots , so $|X| < \min\{\dots, \dots\}$
- If $|X| < \min\{\dots, \dots\}$, then \dots , so $|X| < \text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U})$

Combinatorial characterizations of non(P, QN)

Theorem (Bukovský-Reclaw-Repický, 1991)

$$\text{non}(P, QN) = \text{non}(\text{Fin-P}, \text{Fin-QN}) = \mathfrak{b},$$

where \mathfrak{b} is the **bounding number** i.e. the minimal size of an unbounded family of sequences of natural numbers ordered by the relation \leq^* defined by

$$(a_n) \leq^* (b_n) \iff \exists k \forall n > k (a_n \leq b_n).$$

Definition

Let $\mathfrak{b}_{\mathcal{I}}$ is the minimal size of an unbounded family of sequences of natural numbers ordered by the relation $\leq^{\mathcal{I}}$ defined by

$$(a_n) \leq^{\mathcal{I}} (b_n) \iff \exists A \in \mathcal{I} \forall n \in \mathbb{N} \setminus A (a_n \leq b_n).$$

Remark

In general,

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-QN}) \neq \mathfrak{b}_{\mathcal{I}}!$$

Combinatorial characterization of non(\mathcal{I} -P, \mathcal{I} -QN)

Theorem (F.-Staniszewski, 2015)

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-QN}) = \mathfrak{b}_S(\mathcal{I}),$$

where

$$\mathfrak{b}_S(\mathcal{I}) = \min \left\{ |\mathcal{E}| : \mathcal{E} \subseteq \mathcal{P}_{\mathcal{I}} \wedge \forall \{A_n\} \in \mathcal{P}_{\mathcal{I}} \exists \{E_n\} \in \mathcal{E} \bigcup_{n \in \mathbb{N}} (A_{n+1} \cap \bigcup_{i \leq n} E_i) \notin \mathcal{I} \right\}$$

and $\mathcal{P}_{\mathcal{I}}$ is the family of all partitions $\{A_n : n \in \mathbb{N}\}$ of \mathbb{N} such that $A_n \in \mathcal{I}$ for each $n \in \mathbb{N}$.

Combinatorial characterization of non(\mathcal{I} -P, \mathcal{I} - σ -U)

Theorem (F.-Kwela, 2023)

$$\text{non}(\mathcal{I}\text{-P, } \mathcal{I}\text{-}\sigma\text{-U}) = \mathfrak{b}_\sigma(\mathcal{I}),$$

where

$$\mathfrak{b}_\sigma(\mathcal{I}) = \min\{|\mathcal{E}| : \mathcal{E} \subseteq \mathcal{M}_{\mathcal{I}} \wedge \forall (A_n)_{n \in \mathbb{N}} \in \mathcal{M}_{\mathcal{I}} \exists (E_n)_{n \in \mathbb{N}} \in \mathcal{E} \forall k \exists n > k (E_n \not\subseteq A_n)\}$$

and $\mathcal{M}_{\mathcal{I}}$ is the family of all increasing sequences $(A_n : n \in \mathbb{N})$ such that $A_n \in \mathcal{I}$ for each $n \in \mathbb{N}$.

Combinatorial characterization of non(\mathcal{I} -QN, \mathcal{I} - σ -U)

Theorem (F.-Kwela, 2023)

$$\text{non}(\mathcal{I}\text{-QN}, \mathcal{I}\text{-}\sigma\text{-U}) = \text{add}_\omega(\mathcal{I}),$$

where

$$\text{add}_\omega(\mathcal{I}) = \min\{|\mathcal{A}| : \mathcal{A} \subseteq \mathcal{I} \wedge \forall_{(B_n) \in \mathcal{I}^\omega} \exists_{A \in \mathcal{A}} \forall_n (A \not\subseteq B_n)\}$$

and \mathcal{I}^ω is the family of all sequences $(A_n : n \in \mathbb{N})$ such that $A_n \in \mathcal{I}$ for each $n \in \mathbb{N}$.

Remark

- $\text{add}^*(\mathcal{I}) = \min\{|\mathcal{A}| : \mathcal{A} \subseteq \mathcal{I} \wedge \forall_{B \in \mathcal{I}} \exists_{A \in \mathcal{A}} (A \not\subseteq^* B)\}$
- If \mathcal{I} is a P-ideal, then $\text{add}_\omega(\mathcal{I}) = \text{add}^*(\mathcal{I})$.
- There are non-P-ideals \mathcal{I} with $\text{add}_\omega(\mathcal{I}) \neq \text{add}^*(\mathcal{I}) = \omega$.

Examples

Corollary

$$\mathfrak{b}_\sigma(\mathcal{I}) = \min\{\mathfrak{b}_S(\mathcal{I}), \text{add}_\omega(\mathcal{I})\}$$

Proof

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \min\{\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-QN}), \text{non}(\mathcal{I}\text{-QN}, \mathcal{I}\text{-}\sigma\text{-U})\}$$

Examples

$$\bullet \mathfrak{b}_\sigma(\text{Fin}) = \mathfrak{b}_S(\text{Fin}) = \mathfrak{b} < \infty = \text{add}_\omega(\text{Fin})$$


$$\bullet \mathfrak{b}_\sigma(\mathcal{I}_d) = \text{add}_\omega(\mathcal{I}_d) = \text{add}(\mathcal{N}) \leq \mathfrak{b} = \mathfrak{b}_S(\mathcal{I}_d)$$

$$\bullet \mathfrak{b}_\sigma(\text{Fin} \otimes \text{Fin}) = \mathfrak{b}_S(\text{Fin} \otimes \text{Fin}) = \text{add}_\omega(\text{Fin} \otimes \text{Fin}) = \mathfrak{b}$$

$$\bullet \mathfrak{b}_\sigma(\mathcal{I}_{AD}) = \mathfrak{b}_S(\mathcal{I}_{AD}) = \text{add}_\omega(\mathcal{I}_{AD}) = \omega_1$$

Is distinguishing convergence a topological notion?

Sometimes no!




Theorem

- $\forall (f_n: X \rightarrow \mathbb{R}) \left(f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \iff f_n \xrightarrow{\mathcal{I}\text{-U}} 0 \right) \iff |X| < \omega$
- $\forall (f_n: X \rightarrow \mathbb{R}) \left(f_n \xrightarrow{\mathcal{I}\text{-QN}} 0 \iff f_n \xrightarrow{\mathcal{I}\text{-U}} 0 \right) \iff |X| < \omega$
- $\forall (f_n: X \rightarrow \mathbb{R}) \left(f_n \xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0 \iff f_n \xrightarrow{\mathcal{I}\text{-U}} 0 \right) \iff |X| < \omega$

Is distinguishing convergence a topological notion?

Sometimes yes!



Theorem (F.-Kwela, 2023)

- $|X| < b_S(I) \quad \begin{matrix} \Rightarrow \\ \not\Leftarrow \end{matrix} \quad \forall (f_n: X \rightarrow \mathbb{R}) \quad (f_n \xrightarrow{I-P} 0 \iff f_n \xrightarrow{I-QN} 0)$
- $|X| < b_\sigma(I) \quad \begin{matrix} \Rightarrow \\ \not\Leftarrow \end{matrix} \quad \forall (f_n: X \rightarrow \mathbb{R}) \quad (f_n \xrightarrow{I-P} 0 \iff f_n \xrightarrow{I-\sigma-U} 0)$
- $|X| < \text{add}_\omega(I) \quad \begin{matrix} \Rightarrow \\ \not\Leftarrow \end{matrix} \quad \forall (f_n: X \rightarrow \mathbb{R}) \quad (f_n \xrightarrow{I-QN} 0 \iff f_n \xrightarrow{I-\sigma-U} 0)$

In fact, there exists a Hausdorff compact (hence normal) space X of arbitrary cardinality such that

$$\forall (f_n: X \rightarrow \mathbb{R}) \quad (f_n \xrightarrow{I-P} 0 \iff f_n \xrightarrow{I-\sigma-U} 0)$$

Is distinguishing convergence a topological notion?

Separable spaces

Theorem (F.-Kwela, 2023)

There exists a Hausdorff compact (hence normal) space X of arbitrary cardinality such that

$$\forall_{(f_n: X \rightarrow \mathbb{R})} (f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \iff f_n \xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0)$$

Remark

The above X is a one-point compactification of a discrete topological space of a given cardinality. This space is not separable (unless it is countable) and all but one points are isolated.

Theorem (F.-Kwela, 2023)

There exists a Hausdorff separable, sequentially compact, compact (hence normal) space X of arbitrary cardinality up to \mathfrak{c} such that only countably many points of X are isolated and

$$\forall_{(f_n: X \rightarrow \mathbb{R})} (f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \iff f_n \xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0)$$

Remark

- Since

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \mathfrak{b}_\sigma(\mathcal{I}),$$

there exists a topological space X such that

$$|X| = \mathfrak{b}_\sigma(\mathcal{I}) \quad \text{and} \quad \exists_{(f_n: X \rightarrow \mathbb{R})} (f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \not\iff f_n \xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0)$$

- The space X that is constructed in the proof of the equality

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \mathfrak{b}_\sigma(\mathcal{I})$$

is an uncountable discrete space. In particular, X is not a subspace of reals.

- To obtain X as a subset of reals we need one more characterization of $\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U})$

Subsets of reals distinguishing convergence

Bounding number of a binary relation

Definition

- The set $\mathcal{D}_{\mathcal{I}} \subseteq \mathbb{N}^{\mathbb{N}}$ is defined by

$$x \in \mathcal{D}_{\mathcal{I}} \iff x^{-1}[\{n\}] \in \mathcal{I} \text{ for every } n \in \mathbb{N}$$

- The binary relation $\preceq_{\mathcal{I}}$ on $\mathcal{D}_{\mathcal{I}}$ is defined by

$$x \preceq_{\mathcal{I}} y \iff \{m \in \omega : \exists k \in \omega (x(k) \leq m < y(k))\} \text{ is finite}$$

- The bounding number of the relation \preceq :

$$\begin{aligned} \mathfrak{b}(\preceq_{\mathcal{I}}) &= \min\{|B| : B \text{ is } \preceq_{\mathcal{I}}\text{-unbounded}\} \\ &= \min\{|B| : \forall x \in \mathcal{D}_{\mathcal{I}} \exists y \in B (y \not\preceq_{\mathcal{I}} x)\} \end{aligned}$$

Theorem (F.-Kwela, 2023)

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \mathfrak{b}(\preceq)$$

Subsets of reals distinguishing convergence

Unbounded sets in \preceq

Corollary

$$\mathfrak{b}_\sigma(\mathcal{I}) = \text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \mathfrak{b}(\preceq_{\mathcal{I}})$$

Theorem (F.-Kwela, 2023)

If $X \subseteq \mathbb{N}^{\mathbb{N}}$ is $\preceq_{\mathcal{I}}$ -unbounded, then

$$\exists_{(f_n: X \rightarrow \mathbb{R})} (f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \not\iff f_n \xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0)$$

Corollary

There exists a subset X of **reals** such that $|X| = \mathfrak{b}_\sigma(\mathcal{I})$ and

$$\exists_{(f_n: X \rightarrow \mathbb{R})} (f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \not\iff f_n \xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0)$$

Distinguishing spaces not distinguishing convergence

Corollary

It is consistent that there exists X such that

- $\forall (f_n: X \rightarrow \mathbb{R}) (f_n \xrightarrow{\text{Fin-P}} 0 \iff f_n \xrightarrow{\text{Fin-}\sigma\text{-U}} 0)$
- $\exists (f_n: X \rightarrow \mathbb{R}) (f_n \xrightarrow{\mathcal{I}_d\text{-P}} 0 \not\iff f_n \xrightarrow{\mathcal{I}_d\text{-}\sigma\text{-U}} 0)$

Proof

- $\text{non}(\mathcal{I}_d\text{-P}, \mathcal{I}_d\text{-}\sigma\text{-U}) = \mathfrak{b}_\sigma(\mathcal{I}_d) = \text{add}(\mathcal{N}) \leq \mathfrak{b} = \mathfrak{b}_\sigma(\text{Fin}) = \text{non}(\text{P}, \sigma\text{-U})$
- Consistently: $\text{add}(\mathcal{N}) < \mathfrak{b}$
- Taking X such that $|X| = \text{add}(\mathcal{N})$ and

$$\exists (f_n: X \rightarrow \mathbb{R}) (f_n \xrightarrow{\mathcal{I}_d\text{-P}} 0 \not\iff f_n \xrightarrow{\mathcal{I}_d\text{-}\sigma\text{-U}} 0),$$

we are done.

Distinguishing spaces not distinguishing convergence

Question

Corollary

It is consistent that there exists X such that

- $\forall (f_n: X \rightarrow \mathbb{R}) \left(f_n \xrightarrow{\text{Fin-P}} 0 \iff f_n \xrightarrow{\text{Fin-}\sigma\text{-U}} 0 \right)$
- $\exists (f_n: X \rightarrow \mathbb{R}) \left(f_n \xrightarrow{\mathcal{I}_d\text{-P}} 0 \not\iff f_n \xrightarrow{\mathcal{I}_d\text{-}\sigma\text{-U}} 0 \right)$

Question

Do there exist a space X and an ideal \mathcal{I} such that

- $\exists (f_n: X \rightarrow \mathbb{R}) \left(f_n \xrightarrow{\text{Fin-P}} 0 \not\iff f_n \xrightarrow{\text{Fin-}\sigma\text{-U}} 0 \right)$
- $\forall (f_n: X \rightarrow \mathbb{R}) \left(f_n \xrightarrow{\mathcal{I}\text{-P}} 0 \iff f_n \xrightarrow{\mathcal{I}\text{-}\sigma\text{-U}} 0 \right)$

Remark

- For every ideal \mathcal{I} ,

$$\text{non}(\mathcal{I}\text{-P}, \mathcal{I}\text{-}\sigma\text{-U}) = \mathfrak{b}_\sigma(\mathcal{I}) \leq \mathfrak{b} = \mathfrak{b}_\sigma(\text{Fin}) = \text{non}(\text{P}, \sigma\text{-U})$$

- Hence, the cardinality argument does not work in this case.

- Z. Bukovská, Quasinormal convergence, *Math. Slovaca* 41 (1991), no. 2, 137–146.
- L. Bukovský, I. Reclaw, M. Repický, *Spaces not distinguishing pointwise and quasinormal convergence of real functions*, *Topology Appl.* 41 (1991), no. 1-2, 25–40.
- Á. Császár, M. Laczkovich, Some remarks on discrete Baire classes, *Acta Math. Acad. Sci. Hungar.* 33 (1979), no. 1-2, 51–70.
- R. Filipów, A. Kwela, *Yet another ideal version of the bounding number*, *J. Symb. Log.* 87 (2022), no. 3, 1065–1092
- R. Filipów, M. Staniszewski, *Pointwise versus equal (quasi-normal) convergence via ideals*, *J. Math. Anal. Appl.* 422 (2015), no. 2, 995–1006.